



Advanced Health

Roy Horn DC, FIAMA, PA
9030 W. Fort Island Trail, Suite 10A
Crystal River, Fl. 34429
352-563-2597
www.drhorn.org

Reasons to Eliminate Dairy

April 2008

Milk Substitutes

1 cup = 1 cup of Almond Milk, Rice Milk, Oat Milk

Buttermilk: 1 cup = 1 cup minus 1 tbs. of rice milk or almond milk, plus 1 tbs. lemon juice. Let set for a few minutes.

Butter: 1 tbs. = 1 tbs. sunflower oil or Earth Balance Spread

Creamy Dressing: Mix mayonnaise with your favorite vinaigrette

Heavy Cream: 1 tbs. Tahini dissolved in ¼ cup water (this will not whip)

Think that Dairy doesn't affect you?

Take the challenge: avoid it for 2 weeks and then the next day have all you want.

Couscous and Black Bean Salad

1 Cup couscous; 2 Cups boiling water; 1 ½ C frozen yellow corn; 2 C cooked black beans; 16 roma tomatoes or 3 regular tomatoes, chopped; 8 cloves of garlic, minced; 1 each red and yellow bell pepper, seeded/minced; ¾ C fresh cilantro, minced, optional; 1 ½ tsp salt; 2 tsp cumin; ½ C rice vinegar

Put couscous in a large bowl. Pour the boiling water over it and cover the bowl until the couscous has absorbed all of the water (about 10 mins). Add the corn and mix (the heat from the couscous will thaw the corn). Add the rest of the ingredients and mix well. This tastes great at room temperature, but store in the refrigerator.

Today's Milk

Modern feeding methods substitute high-protein, soy-based feeds for fresh green grass and breeding methods to produce cows with abnormally large pituitary glands so that they produce three times more milk than the old fashioned scrub cow. These cows need antibiotics to keep them well. The pasteurization destroys many valuable enzymes in the milk that are needed to aid digestion. The human pancreas is not always able to produce these enzymes which will over-stress of the pancreas can lead to diabetes and other diseases.

Recombinant Bovine Growth Hormone

Recombinant Bovine Growth Hormone (rBGH) is a genetically engineered, potent variant of the natural growth hormone produced by cows. Injection of this hormone forces cows to increase their milk production by about 10%. rBGH makes cows sick. Monsanto has been forced to admit to about 20 toxic effects, including mastitis, on its POSILAC label. rBGH milk is contaminated by abnormally high pus levels, due to the mastitis commonly induced by rBGH, and antibiotics used to treat the mastitis. rBGH milk is chemically and nutritionally different than natural milk. rBGH milk is contaminated with rBGH, traces of which are absorbed through the gut. rBGH milk is supercharged with high levels of a natural growth factor (IGF-1), which is readily absorbed through the gut. Excess levels of IGF-1 have been incriminated as a cause of breast, colon, and prostate cancers. IGF-1 blocks natural defense mechanisms against early submicroscopic cancers.

Some researchers and "experts" suggest that IGF-1 from outside sources cannot be absorbed because the digestive enzymes destroy it while it's in the GI tract. In 1999, the ADA published research demonstrating that people who consumed 3 servings of milk daily had a 10% higher serum IGF-1 level and almost a 10% lower level IGF Binding Protein 4 (IGBP-4) than those drinking less than 1-1/2 servings.

Constipation

A double-blind trial found that chronic constipation among infants and problems associated with it were triggered by intolerance to cows' milk in two-thirds of the infants studied. Symptoms disappeared in most infants when cows' milk was removed from their diet. *New England Journal of Medicine* 1998;339: pp,1100-4

Dairy Induces Immunization to Insulin

Cow's milk feeding is an environmental trigger of immunity to insulin in infancy that may explain the epidemiological link between the risk of type 1 diabetes and early exposure to cow's milk formulas. *Diabetes*, Vol 48, Issue 7 1389-1394.

High intakes of milk, but not meat, increase serum insulin and insulin resistance in 8-year-old boys.

European Journal of Clinical Nutrition. 2005 Mar;59(3):393-8



The possible role of hormones in milk from cows in the development of breast, ovarian and corpus uteri cancers.

The continued increase in incidence of some hormone-related cancers worldwide is of great concern. Although estrogen-like substances in the environment were blamed for this increase, the possible role of estrogens from food has not been widely discussed. Cows' milk contains a considerable quantity of estrogens. When we name cows' milk as one of the important routes of human exposure to estrogens, the general response of Western people is that "man has been drinking cows' milk for around 2000 years without apparent harm." However, the milk that we are now consuming is quite different from that consumed 100 years ago. Modern dairy cows are usually pregnant and continue to lactate during the latter half of pregnancy, when the concentration of estrogens in blood, and hence in milk, increases. The correlation of incidence and mortality rates with environmental variables in worldwide countries provides useful clues to the etiology of cancer. Among dietary risk factors, we are most concerned with milk and dairy products, because the milk we drink today is produced from pregnant cows, in which estrogen and progesterone levels are markedly elevated. *Medical Hypotheses*. 2005;65(6):1028-37. Epub 2005 Aug 24

Testicular Germ Cell Cancer

Results of a November 2006 case control study suggest that milk fat and/or galactose (a milk sugar) may explain the association between milk and dairy product consumption and seminomatous testicular cancer. *Cancer Epidemiology Biomarkers & Prevention* Vol. 15, 2189-2195, November 2006

Dairy: Colon Cancer Risk Triples

High childhood total dairy intake was associated with a near-tripling in the odds of colorectal cancer in adulthood. *American Journal of Clinical Nutrition*, Vol. 86, No. 6, 1722-1729, December 2007

Ovarian Cancer

Women who consume dairy products on a regular basis, have triple the risk of ovarian cancer than other women. *The Lancet* 1989; 2

In an evaluation of 80,326 women; women who consumed 1+ servings of dairy per day had a 44% greater risk for all types of invasive ovarian cancer compared with those who ate the lowest amount (3 or fewer servings monthly). *American Journal of Epidemiology*, 1999;150

Lymphoma

In Norway, 1422 individuals were followed for 11 ½ years. Those drinking 2 or more glasses of milk per day had 3.5 times the incidence of cancer of the lymphatic organs. *British Med. Journal* 61:456-9, March 1990.

Prostate Cancer

"At least 16 research studies now link milk consumption to prostate cancer, and milk fat is also linked to heart disease," - Neal D. Barnard, M.D. High consumption of dairy products was associated with a 50% increased risk of prostate cancer. *Cancer Causes Control* 1998 Dec;9(6):559-66

Dairy Increases Risk of Squamous Cell Carcinoma in those with History of Skin Cancer

International Journal of Cancer. 2006 Oct 15;119(8):1953-60

Dairy product consumption was positively associated with risk of Parkinson's disease

American Journal of Epidemiology 2007 165(9):998-1006

We believe this is pretty convincing evidence that one should seriously consider eliminating dairy from their diet. Of course, you're probably wondering, "What about my calcium?". Don't worry! We'll answer that question next month.

**Don't Guess About
Your Health...
Schedule a
Nutritional
Consultation
Today!**

563-2597

Our office provides this monthly newsletter free of charge. If you would like to sign up to receive our monthly newsletter and announcements via email, please contact our office

Federal Law requires that we warn you of the following: The information provided in this article is for educational purposes only. Your individual health status and any required health care treatments can only be properly addressed by a professional healthcare provider of your choice. Remember: There is no adequate substitution for a personal consultation with your chosen health care provider. Therefore, we encourage you to make your own health care decisions based upon your research and in partnership with a qualified health care professional. If you are receiving duplicates of this email or would like to be removed from this distribution list, please email Office@DrHorn.org and we can solve the issue. © Copyright Dr. Roy A. Horn. All rights reserved. This content may be copied in full, as long as copyright, contact, and creation information is given, only if used only in a not-for-profit format. If any other use is desired, written permission is required.